


Amendments to the Claims:

The following listing of claims will replace all prior versions, and listings, of claims in the application:

1. (Original) A composition of a ketone peroxide comprising
 - a) a peroxide derivative of the formula
 $\text{HOO-C(R}_1\text{)(R}_2\text{)-OOH}$
wherein
 R_1 is a branched or unbranched alkyl group with 1 to 4 carbon atoms or alkenyl group with 2 to 4 carbon atoms; and
 R_2 is a branched or unbranched alkyl or alkenyl group with 5 to 12 carbon atoms; and
 - b) a branched or unbranched hydrocarbon solvent;
the peroxide derivative of a) having a solubility more than 40 g in 100 g of the solvent of b) at 20°C; and
comprises less than 10 wt.% of a peroxide derivative of the formula
 $\text{HOO-C(R}_1\text{)(R}_2\text{)-OO-C(R}_1\text{)(R}_2\text{)-OOH}$,
wherein R_1 and R_2 have the previously given meanings.
2. (Original) The composition of claim 1 wherein R_1 and R_2 are alkyl groups.
3. (Original) The composition of claim 2 wherein R_1 is a methyl group and R_2 is an isoamyl or amyl group.
4. (Currently Amended) The composition of claim 1 ~~any one of claims 1-3~~ wherein the solvent is a saturated aliphatic hydrocarbon.
5. (Original) A composition of a ketone peroxide derived bis-peroxyester, bis-peroxycarbonate, or mixed peroxyester-peroxycarbonate comprising
 - a) a ketone peroxide derived bis-peroxyester, bis-peroxycarbonate, or mixed peroxyester-peroxycarbonate derivative of the formula



wherein

R_1 is a branched or unbranched alkyl group with 1 to 4 carbon atoms or alkenyl group with 2 to 4 carbon atoms; and

R_2 is a branched or unbranched alkyl or alkenyl group with 5 to 12 carbon atoms; and

R_3 is independently selected from a branched or unbranched alkyl group with 1 to 12 carbon atoms, alkenyl group with 2 to 12 carbon atoms; and an aromatic group with 6-12 carbon atoms,

n is independently 0 or 1, and

b) a branched or unbranched hydrocarbon solvent;

and


comprising less than 10 wt.% of a peroxide derivative of the formula



wherein R_1 , R_2 , R_3 , and n have the previously given meanings.

6. (Original) A composition of a ketone peroxide derived monoperoxyester or monoperoxycarbonate comprising

a) a ketone peroxide derived monoperoxyester or monoperoxycarbonate derivative of the formula



wherein

R_1 is a branched or unbranched alkyl group with 1 to 4 carbon atoms or alkenyl group with 2 to 4 carbon atoms; and

R_2 is a branched or unbranched alkyl or alkenyl group with 5 to 12 carbon atoms; and

R_3 is selected from a branched or unbranched alkyl group with 1 to 12 carbon atoms, alkenyl with 2 to 12 carbon atoms; and an aromatic group with 6-12 carbon atoms;

n is 0 or 1, and

b) a branched or unbranched hydrocarbon solvent;

and

comprising less than 10 wt.% of a peroxide derivative of the formula



wherein R₁, R₂, R₃, and n have the previously given meanings.

7. (Currently Amended) A process for the preparation of a composition of claim 1 ~~any one of the claims 1-4~~ comprising the step wherein a ketone of the formula $\text{O=C(R}_1\text{)(R}_2\text{)}$, wherein R₁ and R₂ have the previously given meanings, is reacted with hydrogen peroxide in the branched or unbranched hydrocarbon solvent in the presence of an acidic catalyst.
8. (Currently Amended) Use of the composition of claim 1 ~~any one of claims 1-6~~ for polymerizing vinylchloride, (meth)acrylic monomers, styrene, ethylene, or mixtures thereof, for curing unsaturated polyester or vinylester resins, for grafting monomers onto a polymer, for crosslinking a polymer or for degrading a polymer.